

Analysis of NO formation during Earth entry using the Quasi Classical Trajectory Method

Narendra Singh

*Department of Aerospace Engineering and Mechanics,
University of Minnesota*

Mentors: Dr. Richard L. Jaffe,
Dr. David W. Schwenke

NASA Ames Research Center

NASA Ames Summer Internship 2017



High Temperature Air Chemistry in Hypersonics

Goal: Understanding high-temperature, non-equilibrium chemistry for hypersonic flows

1. Prediction of the thermo-chemical non-equilibrium state
2. Atomic oxygen and nitrogen dictate heat shield oxidation and ablation
3. Improve thermal protection system design

High Temperature Air Chemistry in Hypersonics

Goal : Understanding high-temperature, non-equilibrium chemistry for hypersonic flows

1. Prediction of the thermo-chemical non-equilibrium state
2. Atomic oxygen and nitrogen dictate heat shield oxidation and ablation
3. Improve thermal protection system design

Existing models :

1. **empirical**
2. **fit to a narrow range of experimental results**
3. **significant uncertainties on extrapolation**

High Temperature Air Chemistry in Hypersonics

Goal : Understanding high-temperature, non-equilibrium chemistry for hypersonic flows

1. Prediction of the thermo-chemical non-equilibrium state
2. Atomic oxygen and nitrogen dictate heat shield oxidation and ablation
3. Improve thermal protection system design

Existing models :

1. **empirical**
2. **fit to a narrow range of experimental results**
3. **significant uncertainties on extrapolation**

Next Generation Models:

1. Based on rigorous framework of statistical thermodynamics
2. Consistent with information from molecular simulations [Quasi Classical Trajectory (QCT) and Direct Molecular Simulation (DMS)]

High Temperature Air Chemistry in Hypersonics

Goal: Understanding high-temperature, non-equilibrium chemistry for hypersonic flows

1. Prediction of the thermo-chemical non-equilibrium state
2. Atomic oxygen and nitrogen dictate heat shield oxidation and ablation
3. Improve thermal protection system design

Existing models :

1. **empirical**
2. **fit to a narrow range of experimental results**
3. **significant uncertainties on extrapolation**

Next Generation Models:


1. Based on rigorous framework of statistical thermodynamics
2. Consistent with information from molecular simulations [Quasi Classical Trajectory (QCT) and Direct Molecular Simulation (DMS)]

Current Focus:

What can we learn from QCT, besides obtaining rate constants, through $\text{N}_2 + \text{O}$ interactions ?


Quasi Classical Trajectory Approach

1. Initialize reactants (quantized)
 - ▶ relative velocity of approach
 - ▶ impact parameter
 - ▶ Internal energy state
 - ▶ Randomize orientations of atoms
2. Integrate Hamilton's equations of motion
3. Analyze outcomes
4. Run lots of these calculations and analyze outcomes (statistically)



Quasi Classical Trajectory Approach

1. Initialize reactants (quantized)
 - ▶ relative velocity of approach
 - ▶ impact parameter
 - ▶ Internal energy state
 - ▶ Randomize orientations of atoms
2. Integrate Hamilton's equations of motion
3. Analyze outcomes
4. Run lots of these calculations and analyze outcomes (statistically)




What do we need for the above? Forces on each atom.

- Potential energy surface (PES)

Quasi Classical Trajectory Approach

1. Initialize reactants (quantized)
 - ▶ relative velocity of approach
 - ▶ impact parameter
 - ▶ Internal energy state
 - ▶ Randomize orientations of atoms
2. Integrate Hamilton's equations of motion
3. Analyze outcomes
4. Run lots of these calculations and analyze outcomes (statistically)



What do we need for the above? Forces on each atom.

- Potential energy surface (PES)

For this work, REAQCT Code, developed at University of Minnesota has been used.

Potential Energy Surface for N_2+O

N_2+O interactions require two surfaces (A'' and A' , ground triplet)




Figure: Contour plot for a possible NON geometry (Energies in kcal/mol)


Lin, W., Varga, Z., Song, G., Paukku, Y. and Truhlar, D.G., 2016. Global triplet potential energy surfaces for the $\text{N}_2(X^1\Sigma)+\text{O}(3P)\rightarrow\text{NO}(X^2\Pi)+\text{N}(4S)$ reaction. The Journal of chemical physics, 144(2), p.024309

NO formation analysis based on molecular energies

Translational energy, $\epsilon_{trans} \approx 1.0$ eV



Dissociation energies: ϵ_d
= 9.91 eV (N_2) and 5.6 eV (NO)



Vibrational energy effect

- linear variation at lower energies
- nearly constant at higher energies


Figure: Exchange reaction probability vs. vibrational energy

NO formation analysis based on molecular energies

Translational energy, $\epsilon_{trans} \approx 1.0$ eV



Dissociation energies: ϵ_d
= 9.91 eV (N_2) and 6.62 eV (NO)



Vibrational energy effect

- linear variation at lower energies
- nearly constant at higher energies


Rotational energy effect

- makes up for threshold energy


Figure: Exchange reaction probability vs. vibrational energy

NO formation analysis based on molecular energies

Translational energy, $\epsilon_{trans} \approx 3.0$ eV



Dissociation energies: ϵ_d
= 9.91 eV (N_2) and 5.6 eV (NO)



Vibrational energy effect

- linear variation at lower energies
- nearly constant at higher energies

Rotational energy effect

- makes up for threshold energy

Translational energy effect


- Increase in ϵ_{trans} increases reaction probability

Figure: Exchange reaction probability vs. vibrational energy


NO formation analysis based on molecular energies

Translational energy, $\epsilon_{trans} \approx 5.0$ eV

Translational energy, $\epsilon_{trans} \approx 8.50$ eV



(a)



(b)

Figure: Exchange reaction probability vs. vibrational energy,
 $N_2(\epsilon_{vib}, \epsilon_{rot}) + O \rightarrow NO + O$

Exchange Reaction Comparison: $\text{N}_2 + \text{O}$ and $\text{N}_2 + \text{N}$ System

Translational energy, $\epsilon_{\text{trans}} \approx 3.00 \text{ eV}$





Figure: Exchange reaction probability vs. vibrational energy,


N₂ dissociation analysis based on molecular energies

Translational energy, $\epsilon_{trans} \approx 1.0$ eV

Translational energy, $\epsilon_{trans} \approx 3.0$ eV



(a)




(b)

Figure: Dissociation reaction probability vs. vibrational energy,
 $\text{N}_2(\epsilon_{vib}, \epsilon_{rot}) + \text{O} \rightarrow \text{N} + \text{N} + \text{O}$


N₂ dissociation analysis based on molecular energies

Translational energy, $\epsilon_{trans} \approx 5.0$ eV



(a)


Translational energy, $\epsilon_{trans} \approx 8.50$ eV



(b)

Figure: Dissociation reaction probability vs. vibrational energy,
 $\text{N}_2(\epsilon_{vib}, \epsilon_{rot}) + \text{O} \rightarrow \text{N} + \text{N} + \text{O}$

Reaction Rate Constants for Various Temperatures



$$k = \frac{1}{3} [k(A'') + k(A')]$$
$$k(A')/k(A'') \approx \frac{1}{2}, T > 12,000K$$
$$\Rightarrow k(\text{approx.}) \approx k(A'')/2$$

Figure: Reaction Rate Constants

Conclusions

- Reaction cross-sections for exchange and dissociation reactions strongly depend on vibrational energy
- Reaction cross-reactions for dissociation has stronger dependence on vibrational energy
- On logarithmic scale, linear variation with vibrational energy for both reactions was shown
- The reaction cross-sections for interactions are not chaotic, in fact they have very simple trends. This is true for non-reactive interactions too (not shown in presentation)
- Simple trends are strong motivation to build simple models consistent with QCT information.

Thank you

Questions?